

## REFERENCES

### A

- Abe, Y., Kude, K., Hayase, S., Kawatsura, M., Tsunashima, K., Itoh, T. (2008). Design of phosphonium ionic liquids for lipase-catalyzed transesterification. *Journal of Molecular Catalysis B: Enzymatic*, 51, 81-5
- Adzamic, T., Sertic-Bionda, K., Zoretic, Z. (2009). Desulfurization of FCC gasoline by extraction with sulfolane and furfural. *NAFTA Journal*, 60, 485-90
- Adzamic, T., Sertic-Bionda, K., Marcec-Rahelic, N. (2010). Modeling of the FCC gasoline desulfurization process by liquid extraction with sulfolane. *Petroleum Science & Technology*, 28, 1936-45
- Alonso<sup>a</sup>, L., Arce, A., Francisco, M., Soto, A. (2007). Liquid-liquid equilibria for systems composed by 1-methyl-3-octylimidazolium tetrafluoroborate ionic liquid, thiophene and *n*-hexane or cyclohexane. *Journal of Chemical & Engineering Data*, 52, 1729-32
- Alonso<sup>b</sup>, L., Arce, A., Francisco, M., Soto, A. (2007). Measurement and correlation of liquid-liquid equilibria of two imidazolium ionic liquids with thiophene and methylcyclohexane. *Journal of Chemical & Engineering Data*, 52, 2409-12
- Alonso<sup>c</sup>, L., Arce, A., Francisco, M., Soto, A. (2007). Gasoline desulfurization using extraction with [C<sub>8</sub>mim][BF<sub>4</sub>] ionic liquid. *AIChE Journal*, 53, 3108-15
- Alonso<sup>a</sup>, L., Arce, A., Francisco, M., Soto, A. (2008). (Liquid + liquid) equilibria of [C<sub>8</sub>mim][NTf<sub>2</sub>] ionic liquid with a sulfur-component and hydrocarbons. *Journal of Chemical Thermodynamics*, 40, 265-70
- Alonso<sup>b</sup>, L., Arce, A., Francisco, M., Soto, A. (2008). Solvent extraction of thiophene from *n*-alkanes (C<sub>7</sub>, C<sub>12</sub> and C<sub>16</sub>) using the ionic liquid [C<sub>8</sub>mim][BF<sub>4</sub>]. *Journal of Chemical Thermodynamics*, 40, 966-72
- Alonso<sup>c</sup>, L., Arce, A., Francisco, M., Soto, A. (2008). Phase behaviour of 1-methyl-3-octylimidazolium bis[trifluoromethylsulfonyl]imide with thiophene and aliphatic hydrocarbons: the influence of *n*-alkane chain length. *Fluid Phase Equilibria*, 263, 176-81
- Alonso<sup>d</sup>, L., Arce, A., Francisco, M., Soto, A. (2008). Thiophene separation from aliphatic hydrocarbons using the 1-ethyl-3-methylimidazolium ethylsulfate ionic liquid. *Fluid Phase Equilibria*, 270, 97-102

Alonso<sup>e</sup>, L., Arce, A., Francisco, M., Soto, A. (2008). Liquid-liquid equilibria of ([C<sub>2</sub>mim][EtSO<sub>4</sub>] + thiophene + 2,2,4-trimethylpentane) and ([C<sub>2</sub>mim][EtSO<sub>4</sub>] + thiophene + toluene): experimental data and correlation. *Journal of Solution Chemistry*, 37, 1355-63

Anantharaj, R. and Banerjee, T. (2011). COSMO-RS based predictions for the desulphurization of diesel oil using ionic liquids: effect of cation and anion combination. *Fuel Processing Technology*, 92, 39-52

Anderson, J.L., Ding, J., Welton, T., Armstrong, D.W. (2002). Characterizing ionic liquids on the basis of multiple solvation interactions. *Journal of American Chemical Society*, 124, 14247-54

Arce<sup>a</sup>, A., Earle, M.J., Katdare, S.P., Rodriguez, H., Seddon, K.R. (2006). Mutually immiscible ionic liquids. *Chemical Communications*, 2548-50

Arce<sup>b</sup>, A., Rodil, E., Soto, A. (2006). Volumetric and viscosity study for the mixtures of 2-ethoxy-2-methylpropane, ethanol and 1-ethyl-3-methylimidazolium ethyl sulfate ionic liquid. *Journal of Chemical & Engineering Data*, 51, 1453-7

Arce<sup>a</sup>, A., Earle, M.J., Katdare, S.P., Rodriguez, H., Seddon, K.R. (2007). Phase equilibria of mixtures of mutually immiscible ionic liquids. *Fluid Phase Equilibria*, 261, 427-33

Arce<sup>b</sup>, A., Earle, M.J., Rodriguez, H., Seddon, K.R. (2007). Separation of benzene and hexane by solvent extraction with 1-alkyl-3-methylimidazolium bis{(trifluoromethyl)sulfonyl}imide ionic liquids: effect of the alkyl-substituent length. *Journal of Physical Chemistry B*, 111, 4732-6

Asian Clean Fuels Association, ACFA (2012). In Focus: Malaysia to implement Euro 4M in 2015. Singapore

## B

Bae, E., Yeo, I.J., Jeong, B., Shin, Y., Shin, K.H., Kim, S. (2011). Study of double bond equivalents and the numbers of carbon and oxygen atom distribution of dissolved organic matter with negative-mode FT-ICR MS. *Analytical Chemistry*, 83, 4193-9

Banerjee, T. and Khanna, A. (2006). Infinite dilution activity coefficients for trihexyltetradecyl phosphonium ionic liquids: measurements and COSMO-RS prediction. *Journal of Chemical & Engineering Data*, 51, 2170-7

Beg, Q., Sahai, V., Gupta, R. (2003). Statistical media optimization and alkine protease production from *Bacillus mojavensis* in bioreactor. *Process Biochemistry*, 39, 203-9

Berthod, A., Ruiz-Angel, M.J., Carda-Broch, S. (2008). Ionic liquids in separation techniques. *Journal of Chromatography A*, 1184, 6-18

Boschetti, A., Montagnaro, F., Rienzo, C., Santoro, L. (2007). A preliminary investigation on the use of organic ionic liquids as green solvents for acylation and oxidation reactions. *Journal of Cleaner Production*, 15, 1797-805

Bosmann, A., Datsevich, L., Jess, A., Lauter, A., Schmitz, C., Wasserscheid, P. (2001). Deep desulfurization of diesel fuel by extraction with ionic liquids. *Chemical Communications*, 2494-5

Brennecke, J.F. and Maginn, E.J. (2001). Ionic liquids: innovative fluids for chemical processing. *AIChE Journal*, 47, 2384-9

Bultinck, P., Ponec, R., Gallegos, A., Fias, S., van Damme, S., Carbo-Dorca, R. (2006). Generalized Polansky index as an aromaticity measure in polycyclic aromatic hydrocarbons. *Croatica Chemica Acta*, 79, 363-71

## C

Cassol, C.C., Umpierre, A.P., Ebeling, G., Ferrera, B., Chiaro, S.S.X., Dupont, J. (2007). On the extraction of aromatic compounds from hydrocarbons by imidazolium ionic liquids. *International Journal of Molecular Sciences*, 8, 593-605

Chen, S.H., Zhao, Q., Xu, X.W. (2008). Preparation and characterization of a novel benzimidazolium Bronsted acidic ionic liquid and its application in esterifications. *Journal of Chemical Science*, 120, 481-3

Chen, X., Li, X., Song, H., Lu, Y., Wang, F., Hu, A. (2008). Synthesis of a basic imidazolide ionic liquid and its application in catalyzing Knoevenagel condensation. *Chinese Journal of Catalysis*, 29, 957-9

Chu, X., Hu, Y., Li, J., Liang, Q., Liu, Y., Zhang, X., Peng, X., Yue, W. (2008). Desulfurization of diesel fuel by extraction with  $[BF_4]^-$ -based ionic liquids. *Chinese Journal of Chemical Engineering*, 16, 881-4

Corma, A., Martinez, C., Ketley, G., Blair, G. (2001). On the mechanism of sulfur removal during catalytic cracking. *Applied Catalyst A: General*, 208, 135-52

Crowhurst, L., Mawdsley, P.R., Perez-Arlandis, J.M., Salter, P.A., Welton, T. (2003). Solvent – solute interactions in ionic liquids. *Physical Chemistry Chemical Physics*, 5, 2790-4

## D

Dai, W., Zhou, Y., Wang, S., Su, W., Sun, Y., Zhou, L. (2008). Desulfurization of transportation fuels targeting at removal of thiophene/benzothiophene. *Fuel Processing Technology*, 89, 749-55

Davila, M.J., Aparicio, S., Alcalde, R., Garcia, B., Leal, J.M. (2007). On the properties of 1-butyl-3-methylimidazolium octylsulfate ionic liquid. *Green Chemistry*, 9, 221-32

Davis, J.H.J. and Forrester, K.J. (1999). Thiazolium-ion based organic ionic liquids (OILs): novel OILs which promote the benzoin condensation. *Tetrahedron Letters*, 40, 1621-2

Deetlefs, M., Seddon, K.R., Shara, M. (2006). Predicting physical properties of ionic liquids. *Physical Chemistry Chemical Physics*, 8, 642-9

Dhir, S., Uppaluri, R., Purkait, M.K. (2009). Oxidative desulfurization: kinetic modeling. *Journal of Hazardous Materials*, 161, 1360-8

Diedenhofen, M., Eckert, F., Klamt, A. (2003). Prediction of infinite dilution activity coefficients of organic compounds in ionic liquids using COSMO-RS. *Journal of Chemical & Engineering Data*, 48, 475-9

Domanska, U. (2005). Solubilities and thermophysical properties of ionic liquids. *Journal of Pure Applied Chemistry*, 77, 543-57

Domanska, U., Pobudkowska, A., Rogalski, M. (2008). Surface tension of binary mixtures of imidazolium and ammonium based ionic liquids with alcohols, or water: cation, anion effect. *Journal of Colloid & Interface Science*, 322, 342-50

Domanska, U. and Krolikowski, M. (2010). Determination of activity coefficients at infinite dilution of 35 solutes in the ionic liquid, 1-butyl-3-methylimidazolium tosylate using gas-liquid chromatography. *Journal of Chemical & Engineering Data*, 55, 4817-22

## E

Earle, M.J. and Seddon, K.R. (2000). Ionic liquids: green solvents for the future. *Journal of Pure Applied Chemistry*, 72, 1391-98

Eßer, J., Wassercheid, P., Jess, A. (2004). Deep desulfurization of oil refinery streams by extraction with ionic liquids. *Green Chemistry*, 6, 316-22

Eckert, F. and Klamt, A. (2002). Fast solvent screening via quantum chemistry: COSMO-RS approach. *AIChE Journal*, 48, 369-85

## G

Gao, H., Luo, M., Xing, J., Wu, Y., Li, Y., Li, W., Liu, Q., Liu, H. (2008). Desulfurization of fuel by extraction with pyridinium-based ionic liquids. *Industrial & Engineering Chemistry Research*, 47, 8384-8

Gao, H., Xing, J., Li, Y., Li, W., Liu, Q., Liu, H. (2009). Desulfurization of diesel fuel by extraction with Lewis-acidic ionic liquids. *Separation Science & Technology*, 44, 971-82

Garcia-Chavez, L.Y., Hermans, A.J., Schuur, B., de Haan, A.B. (2012). COSMO-RS assisted solvent screening for liquid-liquid extraction of mono ethylene glycol from aqueous streams. *Separation & Purification Technology*, 97, 2-10

Garcia-Cruz, I., Valencia, D., Klimova, T., Oviedo-Roa, R., Martinez-Magadan, J.M., Gomez-Balderas, R., Illas, F. (2008). Proton affinity of S-containing aromatic compounds: implications for crude oil hydrodesulfurization. *Journal of Molecular Catalysis A: Chemical*, 281, 79-84

Glasser, L. (2004). Lattice and phase transition thermodynamics of ionic liquids. *Thermochimica Acta*, 421, 87-93

Grensemann, H. and Gmehling, J. (2005). Performance of a conductor-like screening model for real solvents model in comparison to classical group contribution methods. *Industrial & Engineering Chemistry Research*, 44, 1610-24

## H

Hansmeier, A.R., Meindersma, G.W., de Haan, A.B. (2011). Desulfurization and denitrogenation of gasoline and diesel fuels by means of ionic liquids. *Green Chemistry*, 13, 1907-13

Hao, Y., Peng, J., Hu, S., Li, J., Zhai, M. (2010). Thermal decomposition of allyl-imidazolium-based ionic liquid studied by TGA-MS analysis and DFT calculations. *Thermochimica Acta*, 501, 78-83

Heintz, A. (2005). Recent developments in thermodynamics and thermophysics of non-aqueous mixtures containing ionic liquids: a review. *Journal of Chemical Thermodynamics*, 37, 525-35

Heintz, A., Lehmann, J.K., Schmidt, E., Wandschneider, A. (2009). Diffusion coefficients of imidazolium based ionic liquids in aqueous solutions. *Journal of Solution Chemistry*, 38, 1079-83

Holbrey, J.D., Reichert, W.M., Swatloski, R.P., Broker, G.A., Pitner, W.R., Seddon, K.R., Rogers, R.D. (2002). Efficient, halide free synthesis of new, low cost ionic liquids: 1,3-dialkylimidazolium salts containing methyl- and ethyl-sulfate anions. *Green Chemistry*, 4, 407-13

Holbrey, J.D., Reichtart, W.M., Nieuwenhuyzen, M., Sheppard, O., Hardarce, C., Rogers, R.D. (2003). Liquid clathrate formation in ionic liquid – aromatic mixtures. *Chemical Communications*, 476-7

Holbrey, J.D., Martin, I.L., Rothenberg, G., Seddon, K.R., Silvero, G., Zheng, X. (2008). Desulfurisation of oils using ionic liquids: selection of cationic and anionic components to enhance extraction efficiency. *Green Chemistry*, 10, 87-92

Huang, C., Chen, B., Zhang, J., Liu, Z., Li, Y. (2004). Desulfurization of gasoline by extraction with new ionic liquids. *Energy & Fuels*, 18, 1862-4

Huddleston, J.D., Willauer, H.D., Swatloski, R.P., Visser, A.E., Rogers, R.D. (1998). Room temperature ionic liquids as novel media for ‘clean’ liquid-liquid extraction. *Chemical Communications*, 1765-6

Huddleston, J.D., Visser, A.E., Reichert, W.M., Willauer, H.D., Broker, G.A., Rogers, R.D. (2001). Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation. *Green Chemistry*, 3, 156-64

Huo, Y., Xia, S., Ma, P. (2008). Solubility of alcohols and aromatic compounds in imidazolium-based ionic liquids. *Journal of Chemical & Engineering Data*, 53, 2535-9

## I

Ito, E. and van Veen, J.A.R. (2006). On novel processes for removing sulphur from refinery streams. *Catalysis Today*, 116, 446-60

## J

Jiang, X., Nie, Y., Li, C., Wang, Z. (2008). Imidazolium-based alkylphosphate ionic liquids – a potential solvent for extractive desulfurization of fuel. *Fuel*, 87, 79-84

Jin, H., O'Hare, B., Dong, J., Arzhantsev, S., Baker, G.A., Wishart, J.F., Benesi, A.J., Maroncelli, M. (2008). Physical properties of ionic liquids consisting of the 1-butyl-3-methylimidazolium cation with various anions and the bis(trifluoromethylsulfonyl)imide anion with various cations. *Journal of Physical Chemistry*, 112, 81-92

## K

Katritzky, A.R. Karelson, M., Wells, A.P. (1996). Aromaticity as a quantitative concept: aromaticity variation with molecular environment. *Journal of Organic Chemistry*, 61, 1619-23

Kedra-Krolik, K., Fabrice, M., Jaubert, J. N. (2011). Extraction of thiophene or pyridine from *n*-heptane using ionic liquids – gasoline and diesel desulfurization. *Industrial & Engineering Chemistry Research*, 50, 2296-06

Khenkin, A.M. and Neumann, R. (2011). Desulfurization of hydrocarbons by electron transfer oxidative polymerization of heteroaromatic sulfides catalyzed by  $H_5PV_2Mo_{10}O_{40}$  polyoxometalate. *ChemSusChem*, 4, 346-8

Klamt, A. (1995). Conductor-like screening model for real solvents: a new approach to the quantitative calculation of solvation phenomena. *Journal of Physical Chemistry*, 99, 2224-35

Klamt, A. and Eckert, F. (2000). COSMO-RS: a novel and efficient method for the a priori prediction of thermophysical data of liquids. *Fluid Phase Equilibria*, 172, 43-72

Ko, N.H., Lee, J.S., Huh, E.S., Lee, H., Jung, K.D., Kim, H.S., Cheong, M. (2008). Extractive desulfurization using Fe-containing ionic liquids. *Energy & Fuels*, 22, 1687-90

Koch, B.P. and Dittmar, T. (2006). From mass to structure: an aromaticity index for high-resolution mass data of natural organic matter. *Rapid Communications in Mass Spectrometry*, 20, 926-32

Kotelevskii, S.I. and Prezhdo, O.V. (2001). Aromaticity indices revisited: refinement and application to certain five-membered ring heterocycles. *Tetrahedron*, 57, 5715-29

Krossing, I., Slattery, J.M., Daguenet, C., Dyson, P.J., Oleinikova, A., Weingartner, H. (2006). Why are ionic liquids liquid? A simple explanation based on lattice and solvation energies. *Journal of American Chemical Society*, 128, 13427-34

Krygowski, T.M. and Stepien, B.T. (2005). Sigma- and pi-electron delocalization: focus on substituent effects. *Chemical Reviews*, 105, 3482-512

Kuhlmann, E., Haumann, M., Jess, A., Seeberger, A., Wasserscheid, P. (2009). Ionic liquids in refinery desulfurization: Comparison between biphasic and supported ionic liquid phase suspension processes. *ChemSusChem*, 2, 969-77

Kulkarni, P.S. and Afonso, C.A.M. (2010). Deep desulfurization of diesel fuel using ionic liquids: current status and future challenges. *Green Chemistry*, 12, 1139-49

Kumar, A.A.P. and Banerjee, T. (2009). Thiophene separation with ionic liquids for desulphurization: a quantum chemical approach. *Fluid Phase Equilibria*, 278, 1-8

## L

Laus, G., Bentivoglio, G., Schottenberger, H., Kahlenberg, V., Kopacka, H., Roder, T., Sixta, H. (2005). Ionic liquids: current developments, potential and drawbacks for industrial applications. *Lenzinger Berichte*, 84, 71-85

Letcher, T.M., Marciniaik, A., Marciniaik, M., Domanska, U. (2005). Determination of activity coefficients at infinite dilution of solutes in the ionic liquid 1-butyl-3-methylimidazolium octylsulfate using gas-liquid chromatography at a temperature of 298.15K, 313.15K or 328.15K. *Journal of Chemical & Engineering Data*, 50, 1294-8

Li, H., He, L., Lu, J., Zhu, W., Jiang, X., Wang, Y., Yan, Y. (2009). Deep oxidative desulfurization of fuels catalyzed by phosphotungstic acid in ionic liquids at room temperature. *Energy & Fuels*, 23, 1354-7

Likhanova, N.V., Guzman-Lucero, D.G., Flores, E.A., Garcia, P., Dominguez-Aguilar, M., Palomeque, J., Martinez-Palou, R. (2010). Ionic liquids screening for desulfurization of natural gasoline by liquid-liquid extraction. *Molecular Diversity*, 14, 777-87

Liu<sup>a</sup>, S., Xie, C., Yu, S., Liu, F., Ji, K. (2008). Esterification of  $\alpha$ -pinene and acetic acid using acidic ionic liquids as catalysts. *Catalysis Communications*, 9, 1634-8

Liu<sup>b</sup>, D., Gui, J., Song, L., Zhang, X., Sun, Z. (2008). Deep desulfurization of diesel fuel by extraction with task-specific ionic liquids. *Petroleum Science & Technology*, 26, 973-82

Liu, X., Xiaoa, L., Wu, H., Li, Z., Chen, J., Xia, C. (2009). Novel acidic ionic liquids mediated zinc chloride: highly effective catalysts for the Beckmann rearrangement. *Catalysis Communications*, 10, 424-7

Liu, X., Zhou, G., Zhang, X., Zhang, S. (2010). Molecular dynamics simulation of desulfurization by ionic liquids. *AIChE Journal*, 56, 2983-96

Lo, W., Yang, H., Wei, G. (2003). One-pot desulfurization of light oils by chemical oxidation and solvent extraction with room temperature ionic liquids. *Green Chemistry*, 5, 639-42

Lu, L., Cheng, S., Gao, J., Gao, G., He, M. (2007). Deep oxidative desulfurization of fuels catalysed by ionic liquid in the presence of H<sub>2</sub>O<sub>2</sub>. *Energy & Fuels*, 21, 383-4

## M

Ma, X., Sun, L., Song, C. (2002). A new approach to deep desulfurization of gasoline, diesel fuel and jet fuel by selective adsorption for ultra-clean fuels and for fuel cell applications. *Catalysis Today*, 77, 107-16

Marciniak, A. (2011). Influence of anion structure on the liquid-liquid equilibria of 1-ethyl-3-methylimidazolium cation based ionic liquid – hydrocarbon binary systems. *Journal of Chemical & Engineering Data*, 56, 368-74

Marsh, K.N., Deev, A., Wu, A.C-T., Tran, E., Klamt, A. (2002). Room temperature ionic liquids as replacements for conventional solvents – a review. *Korean Journal of Chemical Engineering*, 19, 357-62

Marsh, K.N., Boxall, J.A., Lichtenthaler, R. (2004). Room temperature ionic liquids and their mixtures – a review. *Fluid Phase Equilibria*, 219, 93-8

Matito, E., Duran, M., Sola, M. (2005). The aromatic fluctuation index (FLU): a new aromaticity index based on electron delocalization. *The Journal of Chemical Physics*, 122, 014109

McCabe, W.L., Smith, J.C., Harriott, P. (2001) Unit Operations of Chemical Engineering. 6<sup>th</sup> Edition, United States: McGraw Hill Companies

McCain, W.D. (1990) The Properties of Petroleum Fluids. 2<sup>nd</sup> edition, United States: PennWell Publishing Company

Mei, H., Mei, B.W., Yen, T.F. (2003). A new method for obtaining ultra-low sulfur diesel fuel via ultrasound assisted oxidative desulfurization. *Fuel*, 82, 405-14

Mochizuki, Y. and Sugawara, K. (2008). Removal of organic sulfur from hydrocarbon resources using ionic liquids. *Energy & Fuels*, 22, 3303-7

Mohanty, S., Banerjee, T., Mohanty, K. (2010). Quantum chemical based screening of ionic liquids for the extraction of phenol from aqueous solution. *Industrial & Engineering Chemistry Research*, 49, 2916-25

Montgomery, D.C. (2001) Design & Analysis of Experiments. United States: John Wiley & Sons

Moulijn, J.A., Makkee, M., van Diepen, A. (2001) Chemical Process Technology. United Kingdom: John Wiley & Sons Ltd

## N

Nefedieva, M., Lebedeva, O., Kultin, D., Kustov, L., Borisenkova, S., Krasovskiy, V. (2010). Ionic liquids based on imidazolium tatrafluoroborate for the removal of aromatic sulfur-containing compounds from hydrocarbon mixtures. *Green Chemistry*, 12, 346-9

Nie, Y., Li, C., Sun, A., Meng, H., Wang, Z. (2006). Extractive desulfurization of gasoline using imidazolium-based phosphoric ionic liquids. *Energy & Fuels*, 20, 2083-7

Nie, Y., Li, C., Wang, Z. (2007). Extractive desulfurization of fuel oil using alkylimidazole and its mixture with dialkylphosphate ionic liquids. *Industrial & Engineering Chemistry Research*, 46, 5108-12

Nie, Y., Li, C., Meng, H., Wang, Z. (2008). N,N-dialkylimidazolium dialkylphosphate ionic liquids: their extractive performance for thiophene series compounds from fuel oils versus the length of alkyl group. *Fuel Processing Technology*, 89, 978-83

## O

Oldfield, C., Wood, N.T., Gilbert, S.C., Murray, F.D., Faure, F.R. (1998). Desulphurisation of benzothiophene and dibenzothiophene by actinomycete organisms belonging to the genus *Rhodococcus*, and related taxa. *Antonie van Leeuwenhoek*, 74, 119-32

Oliveira, L.H. and Aznar, M. (2010). Liquid-liquid equilibrium data in ionic liquid + 4-methyldibenzothiophene + *n*-dodecane systems. *Industrial & Engineering Chemistry Research*, 49, 9462-8

Oliveira, L.H. and Aznar, M. (2011). Phase equilibria in ionic liquids + dibenzothiophene + *n*-dodecane systems. *Industrial & Engineering Chemistry Research*, 50, 2289-95

Olivier-Bourbigou, H. and Magna, L. (2002). Ionic liquids: perspectives for organic and catalytic reactions. *Journal of Molecular Catalysis A: Chemical*, 182-183, 419-37

**P**

Palomar, J., Torrecilla, J.S., Ferro, V.R., Rodriguez, F. (2009). Development of an a priori ionic liquid design tool: ionic liquid selection through the prediction of COSMO-RS molecular descriptor by inverse neural network. *Industrial & Engineering Chemistry Research*, 48, 2257-65

Palou, R.M., Likhanova, N.V., Oropeza, E.A.F., Lucero, D.J.G. (2010). Halogen-free ionic liquids in naphtha desulfurization and their recovery. *United States Patent*, US 2010/0051509 A1

Pereiro, A.B., Legido, J.L., Rodriguez, A. (2007). Physical properties of ionic liquids based on 1-alkyl-3-methylimidazolium cation and hexafluorophosphate as anion and temperature dependence. *Journal of Chemical Thermodynamics*, 39, 1168-75

Planeta, J., Karasek, P., Roth, M. (2006). Distribution of sulfur-containing aromatics between [hmim][Tf<sub>2</sub>N] and supercritical CO<sub>2</sub>: a case study for deep desulfurization of oil refinery streams by extraction with ionic liquids. *Green Chemistry*, 8, 70-7

**R**

Raabe, G. and Kohler, J. (2008). Thermodynamical and structural properties of binary mixtures of imidazolium chloride ionic liquids and alcohols from molecular simulation. *Journal of Chemical Physics*, 129, 144503

Rakita, P.E. (2003). Challenges to the commercial production of ionic liquids. *ACS Symposium Series* 856, Chapter 3, 32-40

Rang, H., Kann, J., Oja, V. (2006). Advances in desulfurization research of liquid fuel. *Oil Shale*, 23, 164-76

Revelli, A.L., Mutelet, F., Jaubert, J.N. (2010). Extraction of benzene or thiophene from *n*-heptane using ionic liquids: NMR and thermodynamic study. *Journal of Physical Chemistry B*, 114, 4600-8

Rheinberg, O. V., Lucka, K., Kohne, H., Schade, T., Andersson, J.T. (2008). Selective removal of sulphur in liquid fuels for fuel cell applications. *Fuel*, 87, 2988-96

**S**

Schmidt, R. (2008). [bmim][AlCl<sub>4</sub>] ionic liquid for deep desulfurization of real fuels. *Energy & Fuels*, 22, 1774-8

Schucker, R.C., Baird, W.C. (2001). Electrochemical oxidation of sulfur compounds in naphtha using ionic liquids. *United States Patent*, US 6274026 B1

Shakirullah, M., Ahmad, I., Ahmad, W., Ishaq, M. (2010). Desulphurization study of petroleum products through extraction with aqueous ionic liquids. *Journal of the Chilean Chemical Society*, 55, 179-83

- Shams, S., Bhattacharya, P., Langaas, S., Jacks, G., Ahmed, K.M., Thunkvist, E.L., Chen, D., Granlund, A. (2006). Mapping and interpretation of field data for evaluation and mitigation of groundwater arsenic contamination in Bangladesh. *Journal of Hydroinformatics*, 8, 25-36
- Shariati, A., Gutkowski, K., Peters, C.J. (2005). Comparison of the phase behaviour of some selected binary systems with ionic liquids. *AIChE Journal*, 51, 1532-40
- Shen, B., Li, H., Zhang, W., Zhao, Y., Zhang Z., Wang, X., Shen, S. (2005). A novel composite support for hydrotreating catalyst aimed at ultra-clean fuels. *Catalysis Today*, 106, 206-10
- Shen, C., Li, C.X., Li, X.M., Lu, Y.Z., Muhammad, Y. (2011). Estimation of densities of ionic liquids using Patel-Teja equation of state and critical properties determined from group contribution method. *Chemical Engineering Science*, 66, 2690-8
- Shiflett, M. and Yokozeki, A. (2007). Liqui-liquid equilibria of hydrofluoroethers and ionic liquid 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. *Journal of Chemical & Engineering Data*, 52, 2413-8
- Shiflett, M. and Yokozeki, A. (2008). Liquid-liquid equilibria in binary mixtures containing fluorinated benzenes and ionic liquid 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. *Journal of Chemical & Engineering Data*, 53, 2683-91
- Shiflett, M. and Niehaus, A.M.S. (2010). Liquid-liquid equilibria in binary mixtures containing substituted benzenes with ionic liquid 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. *Journal of Chemical & Engineering Data*, 55, 346-53
- Sidhpuria, K.B. and Parikh, P.A. (2004). Aromatic saturation: a means to cleaner transportation fuels. *Bulletin of the Catalysis Society of India*, 3, 68-71
- Soleimani, M., Bassi, A., Margaritis, A. (2007). Biodesulfurization of refractory organic sulfur compounds in fossil fuels. *Biotechnology Advances*, 25, 570-96
- Song, C. (2003). An overview of new approaches to deep desulfurization for ultra-clean gasoline, diesel fuel and jet fuel. *Catalysis Today*, 86, 211- 63
- Stanislaus, A., Marafi, A., Rana, M.S. (2010). Recent advances in the science and technology of ultra low sulfur diesel (ULSD) production. *Catalysis Today*, 153, 1-68
- Su, B.M., Zhang, S., Zhang, Z.C. (2004). Structural elucidation of thiophene interaction with ionic liquids by multinuclear NMR spectroscopy. *Journal of Physical Chemistry*, 108, 19510-7

Su, W.C., Chou, C.H., Wong, D.S.H., Li, M.H. (2007). Diffusion coefficients and conductivities of alkylimidazolium tetrafluoroborates and hexafluorophosphates. *Fluid Phase Equilibria*, 252, 74-8

## T

Taib, M.M. and Murugesan, T. (2011). Experimental study on the extractive desulfurization of model fuel using hydroxyl ammonium ionic liquids. *Asia-Pacific Journal of Chemical Engineering*

Tariq, M., Forte, P.A.S., Costa Gomes, M.F., Canongia Lopes, J.N., Rebelo, L.P.N. (2009). Densities and refractive indices of imidazolium and phosphonium-based ionic liquids: effect of temperature, alkyl chain length and anion. *Journal of Chemical Thermodynamics*, 41, 790-8

Tian, L., Cao, F., Fang, D., Guo, S. (2007). Alkylation of toluene with 1,3-pentadiene over [bupy]BF<sub>4</sub> – AlCl<sub>3</sub> ionic liquid catalyst. *Chinese Journal in Chemical Engineering*, 15, 680-2

Tokuda, H., Tsuzuki, S., Susan, M.A.H., Hayamizu, K., Watanabe, M. (2006). How ionic are room temperature ionic liquids? An indicator of the physicochemical properties. *Journal of Physical Chemistry*, 110, 19593-600

## V

Visser, A.E., Swatloski, R.P., Reichert, W.M., Griffin, S.T., Rogers, R.D. (2000). Traditional extractants in non-traditional solvents: group1 and 2 extraction by crown ethers in room-temperature ionic liquids. *Industrial & Engineering Chemistry Research*, 39, 3596-604

Visser, A.E., Holbrey, J.D., Rogers, R.D. (2001). Hydrophobic ionic liquids incorporating N-alkylisoquinolinium cations and their utilization in liquid-liquid separations. *Chemical Communications*, 2484-5

## W

Wang<sup>a</sup>, W., Wang, S., Wang, Y., Liu, H., Wang, Z. (2007). A new approach to deep desulfurization of gasoline by electrochemically catalytic oxidation and extraction. *Fuel Processing Technology*, 88, 1002-8

Wang<sup>b</sup>, J., Zhao, D., Zhou, E., Dong, Z. (2007). Desulfurization of gasoline by extraction with N-alkyl-pyridinium-based ionic liquids. *Journal of Fuel Chemistry & Technology*, 35, 293-6

Wang<sup>a</sup>, W., Shao, L., Cheng, W., Yang, J., He, M. (2008). Bronsted acidic ionic liquids as novel catalysts for Prins reaction. *Catalysis Communications*, 9, 337-41

Wang<sup>b</sup>, Y., Jiang, D., Dai, L. (2008). Novel Bronsted acidic ionic liquids based on benzimidazolium cation: synthesis and catalysed acetalization of aromatic aldehydes with diols. *Catalysis Communications*, 9, 2475-80

Wang, X., Han, M., Wan, H., Yang, C., Guan, G. (2011). Study on extraction of thiophene from model gasoline with Bronsted acidic ionic liquids. *Frontiers on Chemical Science & Engineering*, 5, 107-12

Wasserscheid, P., van Hal, R., Bosmann, A. (2002). 1-n-butyl-3-methylimidazolium ([bmim]) octylsulfate – an even ‘greener’ ionic liquid. *Green Chemistry*, 4, 400-4

Wilfred, C.D., Chong, F.K., Man, Z., Bustam, M.A., Abdul Mutualib, M.I., Chan, Z.P. (2012). Extraction of dibenzothiophene from dodecane using ionic liquids. *Fuel Processing Technology*, 93, 85-9

Wilkes, J.S. and Zaworotko, M.J. (1992). Air and water stable 1-ethyl-3-methylimidazolium based ionic liquids. *Journal of the Chemical Society, Chemical Communications*, 13, 965-7

Wong, C.L., Soriano, A.N., Li, M.H. (2009). Infinite dilution diffusion coefficients of [Bmim]-based ionic liquids in water and its molar conductivities. *Journal of Taiwan Institute of Chemical Engineers*, 40, 77-83

Wu, T.Y., Su, S.G., Gung, S.T., Lin, M.W., Lin, Y.C., Lai, C.A., Sun, I.W. (2010). Ionic liquids containing an alkyl sulfate group as potential electrolytes. *Electrochimica Acta*, 55, 4475-82

## Y

Yasuo, K. and Yoshiki, O. (1997). Prospects of petroleum products trades by ROK/Taiwan/Singapore: tougher environmental standards and changes in product imports/exports. *Energy Policy*, 25, 259-65

Yeung, K.Y. and Ruzzo, W.L. (2001). Principal component analysis for clustering gene expression data. *Bioinformatics*, 17, 763-74

Yu, G., Li, X., Liu, X., Asumana, C., Chen, X. (2011). Deep desulfurization of fuel oils using low-viscosity 1-ethyl-3-methylimidazolium dicyanamide ionic liquid. *Industrial & Engineering Chemistry Research*, 50, 2236-44

Yunhai, S., Houyong, S., Deming, L., Qinghua, L., Dexing, C., Yongchuan, Z. (2006). Separation of glycolic acid from glycolonitrile hydrolysate by reactive extraction with tri-n-octylamine. *Separation & Purification Technology*, 49, 20-6

Yunus, N.M., Abdul Mutualib, M.I., Man, Z., Bustam, M.A., Murugesan, T. (2010). Thermophysical properties of 1-alkylpyridinium bis(trifluoromethylsulfonyl)imide ionic liquids. *Journal of Chemical Thermodynamics*, 42, 491-5

## Z

Zhang, S. and Zhang, Z.C. (2002). Novel properties of ionic liquids in selective sulfur removal from fuels at room temperature. *Green Chemistry*, 4, 376-9

Zhang, S., Zhang, Q., Zhang, Z.C. (2004). Extractive desulfurization and denitrogenation of fuels using ionic liquids. *Industrial & Engineering Chemistry Research*, 43, 614-22

Zhao, D., Wang, J., Zhou, E. (2007). Oxidative desulfurization of diesel fuel using Bronsted acid room temperature ionic liquid in the presence of H<sub>2</sub>O<sub>2</sub>. *Green Chemistry*, 9, 1219-22.

Zhao, D., Sun, Z., Li, F., Liu, R., Shan, H. (2008). Oxidative desulfurization of thiophene catalyzed by (C<sub>4</sub>H<sub>9</sub>)<sub>4</sub>NBr·2C<sub>6</sub>H<sub>11</sub>NO coordinated ionic liquid. *Energy & Fuels*, 22, 3065-9

Zhou, J., Mao, J., Zhang, S. (2008). Ab initio calculations of the interaction between thiophene and ionic liquids. *Fuel Processing Technology*, 89, 1456-60